

AD-A150 112

RECURSIVE MOMENT FORMULAS FOR REGENERATIVE SIMULATION

1/1

(U) STANFORD UNIV CA DEPT OF OPERATIONS RESEARCH

P W GLYNN ET AL. OCT 84 TR-5 ARO-20927.5-MA

UNCLASSIFIED

DAG29-84-K-0030


F/G 12/1

NL

END

PAVED

ATA



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

ARO 20927.5-MA  
②

AD-A150 112

RECURSIVE MOMENT FORMULAS FOR REGENERATIVE SIMULATION

by

Peter W. Glynn and Donald L. Iglehart

TECHNICAL REPORT NO. 5

October 1984

Prepared under the Auspices


of

U.S. Army Research Office Contract\*  
DAAG29-84-K-0030

Approved for public release: distribution unlimited.

Reproduction in Whole or in Part is Permitted for any  
Purpose of the United States Government.

DEPARTMENT OF OPERATIONS RESEARCH  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA



\*This research was also partially supported under  
National Science Foundation Grant MCS-8203483

DTIC FILE COPY

85 01 28 086

# RECURSIVE MOMENT FORMULAS FOR REGENERATIVE SIMULATION

by

Peter W. Glynn  
University of Wisconsin

and

Donald L. Iglehart  
Stanford University

## 1. Introduction

Let  $f$  be a real-valued function defined on the state space of a regenerative process  $X = \{X(t) : t \geq 0\}$  with regeneration times  $0 = T_0 < T_1 < \dots$ , and suppose that

$$(1.1) \quad r_t = \frac{1}{t} \int_0^t f(X(s)) ds \rightarrow r \quad \text{a.s.}$$

as  $t \rightarrow \infty$ . The problem of estimating  $r$  via a simulation of  $X$  is called the steady state simulation problem.

Relation (1.1) implies that  $r_t$  is a strongly consistent point estimator for  $r$ . To obtain confidence intervals for  $r$ , set (for  $k \geq 1$ )

$$Y_k(f) = \int_{T_{k-1}}^{T_k} f(X(s)) ds$$

$$\tau_k = T_k - T_{k-1}$$

$$Z_k = Y_k(f) - r\tau_k.$$

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>
By _____	
Distribution _____	
Availability Codes	
Dist _____	
A-1	

The regenerative structure of  $\mathbf{X}$  guarantees that  $r = E\{Y_1(f)\}/E\{\tau_1\}$  and that  $\{(Y_k(f), \tau_k) : k \geq 1\}$  is a sequence of i.i.d. random vectors. Standard arguments (see CRANE and IGLEHART (1975)) show that if  $E\{Y_1^2(f) + \tau_1^2\} < \infty$ , then

$$(1.2) \quad \sqrt{t}(r_t - r) \Rightarrow \sigma N(0,1)$$

as  $t \rightarrow \infty$ , where  $\sigma^2 = \sigma^2\{Z_1\}/E\{\tau_1\}$ . To use (1.2) for confidence intervals, the regenerative cycle structure of  $\mathbf{X}$  is exploited to obtain a strongly consistent estimator  $v_t$  for  $\sigma^2$ .

These confidence intervals, while asymptotically correct, often have poor small-sample behavior. For example, such confidence intervals often tend to significantly undercover the parameter  $r$ . Several recent studies have examined this problem. GLYNN (1982), in considering regenerative confidence intervals on the time scale of regenerative cycles, obtained asymptotic expansions for the coverage error which indicated that skewness/kurtosis effects play a significant role in determining quality of the confidence interval. To be more precise, the error, to a first approximation, is determined by the magnitude of quantities of the form  $E\{Z_1^m \tau_1^n\}$  for  $m + n \leq 4$ . GLYNN and IGLEHART (1984) obtain expressions for the asymptotic covariance between  $r_t$  and  $v_t$ , and the variance of  $v_t$ ; these expressions also involve mixed moments of the form  $E\{Z_1^m \tau_1^n\}$ . Consequently, in studying small-sample behavior of regenerative confidence intervals, it is of some interest to be able to calculate the exact values of the mixed moments for some "test case" stochastic models. HORDIJK,

IGLEHART, and SCHAASSBERGER (1976) showed how to do this for  $m + n \leq 2$ , when  $X$  is a discrete or continuous time Markov chain with countably many states. In this note, we show how to calculate such quantities when  $X$  is a semi-Markov process with countably many states; discrete and continuous time Markov chain results follow as special cases.

## 2. Statement of the Recursive Moment Formulas

Let  $X = \{X(t) : t \geq 0\}$  be an irreducible non-explosive regenerative semi-Markov process on countable state space  $E$ . Thus,  $X(t)$  may be represented as

$$X(t) = \sum_{k=0}^{\infty} R_k I(S_k \leq t < S_{k+1}) ,$$

where:

- (i)  $R = \{R_n : n \geq 0\}$  is a discrete-time Markov chain on  $E$  with transition matrix  $P = (p_{xy} : x, y \in E)$
- (ii)  $S = \{S_n : n \geq 0\}$  is an increasing sequence of jump times with  $S_0 = 0$  and differences  $\alpha_n = S_{n+1} - S_n$  which are conditionally independent r.v.'s given  $R$ .

The conditional distribution of  $\alpha_n$  is given by  $F(R_n, R_{n+1}, dt) = P(\alpha_n \in dt | R)$ , where  $F(x, y, 0) = 0$  for all  $x, y \in E$ . Note that  $S_n \rightarrow \infty$  a.s., since  $X$  is non-explosive by assumption. Fix  $z \in E$  as the regenerative state; let  $T(z) = \inf\{t > 0 : X(t-) \neq z, X(t) = z\}$  and set

$$Y(u) = \int_0^{T(z)} u(X(t)) dt$$

where  $u : E \rightarrow \mathbb{R}$  is an arbitrary function. We wish to study mixed moments of the form

$$a_{ij}(x) = E_x \{ Y(g)^i Y(h)^j \}$$

for  $x \in E$ ,  $0 \leq i \leq m$ , and  $0 \leq j \leq n$ , when  $g$  and  $h$  are fixed functions, and  $m$  and  $n$  are non-negative integers. Throughout the paper we shall use  $P_x\{\cdot\}$  and  $E_x\{\cdot\}$  to denote conditional probabilities and expectations, given  $X(0) = R_0 = x$ . Note that by choosing  $g(\cdot) = f(\cdot) - r$  and  $h(\cdot) = 1$ ,  $a_{ij}(z)$  yields  $E_z\{Z_1^i \tau_1^j\}$ .

To state our result, let  $a \vee b$  denote  $\max(a, b)$  and set  $b_{mn}(x) = E_x\{((Y(|g|) \vee 1)^m (Y(|h|) \vee 1)^n)\}$ . Let  $G_n$  and  $\beta_n$  be the matrix and function, respectively, defined by

$$G_n(x, y) = \begin{cases} p_{xy} \mu_n(x, y) & ; \quad y \neq z \\ 0 & ; \quad y = z \end{cases}$$

$$\beta_n(x) = \sum_{y \in E} \mu_n(x, y) p_{xy}$$

where  $\mu_n(x, y) = \int_0^\infty t^n F(x, y, dt)$ . Also, we shall identify real-valued functions  $u(\cdot)$  on  $E$  with column vectors  $u$ , and shall use the notation  $u \circ v$  to denote the vector with  $x^{\text{th}}$  component  $(u \circ v)(x) = u(x)v(x)$ . Set  $u^0(\cdot) = 1$  and  $u^{n+1} = (u \circ u^n)$  for  $n \geq 0$ .

Let  $\mathcal{C}$  denote the class of all  $m \times n$  matrix-valued functions,  $\mathcal{C}$ , on  $E$ . Then set

$$_{mn} = \{C \in C : \sum_{k=0}^i \sum_{l=0}^j |g(x)|^{i-k} \cdot |h(x)|^{j-l}$$

$$\cdot \sum_{y \in E} G_{i+j-k-l}(x,y) |c_{kl}(y)| < \infty$$

$$\text{and } (G_0^k c_{ij})(x) \rightarrow 0 \text{ as } k \rightarrow \infty,$$

$$\text{for all } x \in E, 0 \leq i \leq m, \text{ and } 0 \leq j \leq n\}.$$

(2.1) Theorem. If  $b_{mn}(z) < \infty$ , then  $A = \{a_{ij} : 0 \leq i \leq m, 0 \leq j \leq n\}$  is the unique solution in  $C_{mn}$  to the system:

$$(2.2) c_{ij} = g^j \circ h^j \circ \beta_{i+j} + \sum_{(k,l) \in B_{ij}} \binom{i}{k} \binom{j}{l} (g^{i-k} \circ h^{j-l} \circ G^{i+j-k-l} c_{kl}),$$

where  $0 \leq i \leq m, 0 \leq j \leq n$ , and  $B_{ij} = \{(k,l) : 0 \leq k \leq i, 0 \leq l \leq j, k+l > 0\}$ .

Set  $\tau = \inf\{n \geq 1 : R_n = z\}$  and observe that

$$G_0^k(x,y) = P_x\{R_k = y, \tau > k\}.$$

Hence,  $G_0^k \rightarrow 0$  as  $k \rightarrow \infty$ . It follows that if  $E$  has a finite number of elements,  $C_{mn} \equiv C$ , so that the  $a_{ij}$ 's are unique in the class of all possible solutions to (2.2). Also, in the presence of a finite state space, it is well known that  $(I - G_0)^{-1}$  exists, so that (2.2) may be re-written as



$$(2.3) \quad c_{ij} = (I - G_0)^{-1} \{ g^i \circ h^j \circ \beta_{i+j} \\ + \sum_{(k,l) \in A_{ij}} \binom{i}{k} \binom{j}{l} (g^{i-k} \circ h^{j-l} \circ (G_{i+j-k-l} c_{kl})) \} ,$$

where  $0 \leq i \leq m$ ,  $0 \leq j \leq n$ ,  $i+j > 0$ , and  $A_{ij} = \{(k,l) : 0 \leq k \leq i, 0 \leq l \leq j, 0 < k+l < i+j\}$ . Also observe that  $c_{00} \equiv 1$ . Note that the system of equations (2.3) is recursive in  $i+j$ , in the sense that the  $c_{ij}$ 's may be solved in terms of the  $c_{kl}$ 's, where  $k+l < i+j$ . By successively solving for the  $c_{kl}$ 's with fixed  $k+l$  on each iteration, one eventually obtains  $c_{mn}$ .

Formula (2.3) can be further simplified when  $X$  has special structure. Note that if  $X$  is a continuous time Markov chain, then

$$F(x,y,dt) = \lambda(x) \exp(-\lambda(x)t) dt$$

for  $t > 0$ , so that

$$\mu_n(x,y) = n!/\lambda(x)^n \equiv \eta_n(x) .$$

We find that (2.3) can be re-written as

$$c_{ij} = (I - G_0)^{-1} \{ g^i \circ h^j \circ \eta_{i+j} \\ + \sum_{(k,l) \in A_{ij}} \binom{i}{k} \binom{j}{l} (g^{i-k} \circ h^{j-l} \circ \eta_{i+j-k-l} \circ G_0 c_{kl}) \} ,$$

where  $0 \leq i \leq m$ ,  $0 \leq j \leq n$ , and  $i+j > 0$ .

For discrete time Markov chains,  $\beta_n \equiv 1$  and  $G_n = G_0$ , so (2.3) takes the form

$$c_{ij} = (I - G_0)^{-1} \{ g^i \circ h^j + \sum_{(k,l) \in A_{ij}} \binom{i}{k} \binom{j}{l} (g^{i-k} \circ h^{j-l} \circ (G_0 c_{kl})) \} ,$$

where  $0 \leq i \leq m$ ,  $0 \leq j \leq m$ , and  $i+j > 0$ .

Relation (2.4) expresses  $c_{ij}$  in terms of  $G_0 c_{kl}$ , where  $(k,l) \in A_{ij}$ . Equation (2.4) can be re-written, when  $g = h$ , so that the  $c_{ij}$ 's are written directly in terms of the  $c_{kl}$ 's. If  $g = h$ , we write  $c_{ij}$  as  $c_{i+j}$ , and observe that (2.4) takes the form

$$(2.5) \quad c_i = (I - G_0)^{-1} \{ g^i + \sum_{k=1}^{i-1} \binom{i}{k} (g^{i-k} \circ G_0 c_k) \}, \quad 1 \leq i \leq n.$$

Recall also that from (2.2),  $c_0 \equiv 1$ . We claim that the system (2.5) can be re-written as

$$(2.6) \quad c_i = (I - G_0)^{-1} \left\{ \sum_{k=1}^i (-1)^{k+1} \binom{i}{k} g^k \circ c_{i-k} \right\}, \quad 1 \leq i \leq n.$$

The proof is by induction. For  $n = 1$ , the result is obvious, so suppose (2.5) and (2.6) are equivalent systems for  $n = m$ . To check the  $(m+1)$ 'st equation in the  $(m+1)$ 'st system, observe that a solution of (2.5) satisfies

$$(2.7) \quad c_{m+1} = (I - G_0)^{-1} \left\{ g^{m+1} + \sum_{i=1}^m \binom{m+1}{i} (g^{m+1-i} \circ G_0 c_i) \right\}.$$

By the inductive hypothesis, (2.6) shows that

$$(2.8) \quad G_0 c_i = - \sum_{k=0}^i (-1)^{k+1} \binom{i}{k} g^k \circ c_{i-k}$$

for  $i \leq m$ . Substituting (2.8) into (2.7), we get that  $(I - G_0)c_{m+1}$

equals

$$\begin{aligned} g^{m+1} + \sum_{i=1}^m \sum_{k=0}^i \binom{m+1}{i} \binom{i}{k} (-1)^{i-k} g^{m+1-k} \circ c_k \\ = \sum_{k=0}^m \binom{m+1}{k} (g^{m+1-k} \circ c_k) \sum_{j=1}^{m+1-k} \binom{m+1-k}{j} (-1)^{m+1-k-j} \\ = \sum_{k=0}^m \binom{m+1}{k} (g^{m+1-k} \circ c_k) (-1)^{m+2-k} \\ = \sum_{l=1}^{m+1} \binom{m+1}{l} (g^l \circ c_{m+1-l}) (-1)^{l+1}, \end{aligned}$$

which is equivalent to the  $(m+1)$ 'st relation of (2.6) (the binomial identity was used for the third equality). The steps being reversible, this proves the claimed result. We remark that (2.6) yields the equations of [4] for  $i = 1, 2$ .

### 3. Proof of the Theorem

We proceed via a series of lemmas.

(3.1) **Lemma.** If  $b_{mn}(z) < \infty$ , then the r.v.'s  $Y(g)^i Y(h)^j$  are integrable under the probability distribution  $P_x$  for  $0 \leq i \leq m$ ,  $0 \leq j \leq n$ ,  $x \in E$ .

**Proof.** Note that

$$(Y(|g|) \vee 1)^i (Y(|h|) \vee 1)^j \leq (Y(|g|) \vee 1)^m (Y(|h|) \vee 1)^n$$

so that  $b_{ij}(z) < \infty$  for  $0 \leq i \leq m, 0 \leq j \leq n$ . Since  $X$  is irreducible, it follows that  $P_z\{T(y) < T(z)\} > 0$  for all  $y \in E$ , for otherwise the regenerative property guarantees that  $P_z\{T(y) = \infty\} = 1$ , which violates our irreducibility assumption.

Then, by the strong Markov property applied at time  $T(y)$ ,

$$\begin{aligned} E_z \left\{ \left( \int_{T(y)}^{T(z)} |g(X(s))| ds \vee 1 \right)^i \left( \int_{T(y)}^{T(z)} |h(X(s))| ds \vee 1 \right)^j; T(y) < T(z) \right\} \\ = b_{ij}(y) P\{T(y) < T(z) | X(0) = z\} \leq b_{ij}(z) \end{aligned}$$

so that  $b_{ij}(y) < \infty$  for all  $y \in E$ , which proves the result.  $\square$

**(3.2) Proposition.** If  $b_{mn}(z) < \infty$ , then  $a_{ij}(x)$  exists and is finite for  $0 \leq i \leq m, 0 \leq j \leq n, x \in E$ . Furthermore,  $A$  solves (2.2).

**Proof.** The first part follows immediately from Lemma 3.1. For the second part, the integrability of  $Y(g)^i Y(h)^j$  ensures that the following manipulations of conditional expectations are valid:

$$\begin{aligned}
a_{ij}(x) &= E_x \{ Y(g)^i Y(h)^j \} \\
&= E_x \{ Y(g)^i Y(h)^j; \tau = 1 \} + E_x \{ Y(g)^i Y(h)^j; \tau > 1 \} \\
&= g^i(x) h^j(x) \mu_{i+j}(x, z) P_{xz} \\
&\quad + E_x \left\{ (g(R_0) \alpha_0 + \int_{S_1}^{T(z)} g(X(t)) dt)^i (h(R_0) \alpha_0 \right. \\
&\quad \left. + \int_{S_1}^{T(z)} h(X(t)) dt)^j; \tau > 1 \right\} \\
&= g^i(x) h^j(x) \mu_{i+j}(x, z) P_{xz} \\
&\quad + \sum_{k=0}^i \sum_{l=0}^j \binom{i}{k} \binom{j}{l} E_x \{ g(R_0)^{i-k} h(R_0)^{j-l} \alpha_0^{i+j-k-l} \left( \int_{S_1}^{T(z)} g(X(t)) dt \right)^k \\
&\quad \cdot \left( \int_{S_1}^{T(z)} h(X(t)) dt \right)^l; \tau > 1 \} \\
&= g^i(x) h^j(x) \mu_{i+j}(x, z) P_{xz} \\
&\quad + \sum_{k=0}^i \sum_{l=0}^j \binom{i}{k} \binom{j}{l} g^{i-k}(x) h^{j-l}(x) \sum_{y \neq z} P_{xy} \mu_{i+j-k-l}(x, y) a_{kl}(y) \\
&= (g^i \circ h^j \circ \beta_{i+j})(x) + \sum_{(k,l) \in B_{ij}} \binom{i}{k} \binom{j}{l} (g^{i-k} \circ h^{j-l} \\
&\quad \circ (G_{i+j-k-l} a_{kl}))(x) ;
\end{aligned}$$

the strong Markov property at time  $S_1$  was used to obtain the second last equality.  $\square$

(3.3) **Lemma.** If  $b_{mn}(z) < \infty$ , then  $A \in C_{mn}$ .

**Proof.** For the absolute summability observe that  $d_{ij}(x) = E_x\{Y(|g|)^i Y(|h|)^j\}$  satisfies

$$d_{ij} = |g|^i \circ |h|^j \circ \beta_{i+j} + \sum_{(k,l) \in B_{ij}} \binom{i}{k} \binom{j}{l} (|g|^{i-k} \circ |h|^{j-l} \circ (G_{i+j-k-l} d_{kl})) .$$

But  $d_{ij}(x) \leq b_{ij}(x) < \infty$ , so  $|g|^{i-k} \circ |h|^{j-l} \circ (G_{i+j-k-l} d_{kl})$  is finite, proving the first part, since  $|a_{ij}| \leq d_{ij}$ . For the second,

$$(G_0^k a_{ij})(x) = E_x \left\{ \left( \int_{S_k}^{T(z)} Y(g) \right)^i \left( \int_{S_k}^{T(z)} Y(h) \right)^j ; \tau > k \right\} ,$$

which tends to zero by integrability of  $Y(|g|)^i Y(|h|)^j$ .

**(3.4) Lemma.** If  $b_{mn}(z) < \infty$ , then  $A$  is the unique solution to (2.2) in  $C_{mn}$ .

**Proof.** Suppose that  $\{a_{rs} : 0 \leq r \leq k, 0 \leq s \leq l\}$  is unique in  $C_{kl}$  for  $k+l < i+j$ , where  $0 \leq i \leq m, 0 \leq j \leq n$ . We shall prove uniqueness in  $C_{ij}$ ; "bootstrapping" this result yields the lemma. Since the  $a_{rs}$ 's are unique in  $C_{kl}$ , any solution to the  $(i,j)$  equation must satisfy

$$(3.5) \quad c_{ij} = |g|^i \circ |h|^j \circ \beta_{i+j} + \sum_{(k,l) \in A_{ij}} \binom{i}{k} \binom{j}{l} (|g|^{i-k} \circ |h|^{j-l} \circ c_{i+j-k-l} a_{kl}) + G_0 c_{ij} .$$

By the absolute summability, any two solutions  $c_{ij}, c'_{ij}$  of (3.5) must satisfy  $(c_{ij} - c'_{ij}) = G_0(c_{ij} - c'_{ij})$ . Since  $G_0^k(c_{ij} - c'_{ij}) > 0$  as  $k \rightarrow \infty$ , it follows that  $c_{ij} = c'_{ij}$ , proving uniqueness in  $C_{ij}$ .  $\square$

These above results prove all the assertions of the theorem.

#### **Acknowledgment**

Both authors gratefully acknowledge support by U.S. Army Research Office Contract DAAG29-84-K-0030. The second author was also partially supported by National Science Foundation Grant MCS-8203483.

**References:**

- [1] CRANE, M.A. and IGLEHART, D.L. (1975). "Simulating Stable Stochastic Systems, III: Regenerative Processes and Discrete-Event Simulations," Oper. Res. 23, 33-45.
- [2] GLYNN, P.W. (1982). "Asymptotic Theory for Nonparametric Confidence Intervals," Technical Report No. 19. Dept. of Operations Research, Stanford University, Stanford, CA.
- [3] GLYNN, P.W. and IGLEHART, D.L. (1984). "The Joint Limit Distribution of the Sample Mean and the Regenerative Variance Estimator." Forthcoming technical report, Dept. of Operations Research, Stanford University, Stanford, CA.
- [4] HORDIJK, A., IGLEHART, D.L., and SCHASSBERGER, R. (1976). "Discrete-Time Methods of Simulating Continuous-Time Markov Chains," Advances in Appl. Probability 8, 772-788.



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 5	2. GOVT ACCESSION NO. N/A	3. RECIPIENT'S CATALOG NUMBER N/A
4. TITLE (and Subtitle)  RECURSIVE MOMENT FORMULAS FOR REGENERATIVE SIMULATION		5. TYPE OF REPORT & PERIOD COVERED  TECHNICAL REPORT
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) PETER W. GLYNN DONALD L. IGLEHART		8. CONTRACT OR GRANT NUMBER(s)  DAAG29-84-K-0030
9. PERFORMING ORGANIZATION NAME AND ADDRESS DEPARTMENT OF OPERATIONS RESEARCH STANFORD UNIVERSITY STANFORD, CA 94305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709		12. REPORT DATE OCTOBER 1984
		13. NUMBER OF PAGES 13
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)  Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  NA		
18. SUPPLEMENTARY NOTES  The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  REGENERATIVE SIMULATION, SEMI-MARKOV PROCESS, RECURSIVE MOMENT FORMULAS.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  an expression ABSTRACT: Let $\{X(t) : t \geq 0\}$ be a regenerative semi-Markov process with countable state space $E$ and $f$ a real-valued function on $E$ . Denote by $Y_1(f)$ the area under the function $f(X(\cdot))$ in the first regenerative cycle. This paper gives a recursive method for computing moments of the form $E\{Y_1^m(f)Y_1^n(g)\}$ for arbitrary $f$ and $g$ , and $m, n \geq 1$ . These moments are needed to improve the accuracy of confidence intervals for steady state parameters obtained when using the method of regenerative simulation.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

**END**

**FILMED**

**3-85**

**DTIC**